

Nonlinear integral equations for thermodynamics of the $sl(r+1)$ Uimin-Sutherland model

Zengo Tsuboi *

*Graduate School of Mathematical Sciences, University of Tokyo,
Komaba 3-8-1, Meguro-ku, Tokyo 153-8914, Japan*

Abstract

We derive traditional thermodynamic Bethe ansatz (TBA) equations for the $sl(r+1)$ Uimin-Sutherland model from the T -system of the quantum transfer matrix. These TBA equations are identical to the ones from the string hypothesis. Next we derive a new family of nonlinear integral equations (NLIE). In particular, a subset of these NLIE forms a system of NLIE which contains only a *finite* number of unknown functions. For $r = 1$, this subset of NLIE reduces to Takahashi's NLIE for the XXX spin chain. A relation between the traditional TBA equations and our new NLIE is clarified. Based on our new NLIE, we also calculate the high temperature expansion of the free energy.

Short title: Nonlinear integral equations

PACS2001: 02.30.Rz; 02.30.Ik; 05.50.+q; 05.70.-a

MSC: 82B23; 45G15; 82B20; 17B80

Key words: nonlinear integral equation; Uimin-Sutherland model; quantum transfer matrix; solvable lattice model; thermodynamic Bethe ansatz; T -system

to appear in J. Phys. A: Math. Gen.

*E-mail address: ztsuboi@poisson.ms.u-tokyo.ac.jp

1 Introduction

Thermodynamic Bethe ansatz (TBA) equations have been used to investigate thermodynamics of various kind of solvable lattice models (see for example [1]). Traditionally, TBA equations have been derived by the string-hypothesis [2, 3]. Several years ago, TBA equations for the supersymmetric $t - J$ model and the supersymmetric extended Hubbard model were derived [4] from the T -system (a system of functional relations among transfer matrices) for the quantum transfer matrix (QTM) [5, 6, 7, 8, 9], which is independent of the string hypothesis. Later on, TBA equations for the XXZ -model in the regime $|\Delta| < 1$ [10], the $osp(1|2)$ model [11], the $osp(1|2s)$ model [12] were derived by the similar procedures in [4]. In addition, the TBA equation in [10] was analytically continued [13] to the one for $|\Delta| \geq 1$. These TBA equations contain an infinite (or finitely many) number of unknown functions, and thus are not always easily treated. It is significant to simplify the TBA equations to tractable integral equations which contain only a finite number of unknown functions.

In the context of the QTM method, for models related to algebras of rank one or two, nonlinear integral equations (NLIE) with finite numbers of unknown functions were derived by Klümper and his collaborators [9, 14, 15, 16, 17, 18]. Although their NLIE give the same free energy as TBA equations, the derivations need trial and error for each model, which prevents their extension to higher rank case.

Another type of NLIE with only one unknown function was proposed for the XXZ spin chain by Takahashi [19] recently. One can also derive [13] Takahashi's NLIE from the T -system [20, 21] of the QTM. In view of this fact, we have derived [22] NLIE with a finite number of unknown functions from our T -system [23] for the $osp(1|2s)$ model for arbitrary rank s . In this paper, we shall further derive NLIE for the $sl(r+1)$ Uimin-Sutherland model [24, 25] with only a *finite* number (the number of rank r) of unknown functions from the T -system [21]. This is the first explicit derivation of this type of NLIE for a vertex model associated with $sl(r+1)$ for *arbitrary* rank r .

In section 2, we introduce the $sl(r+1)$ Uimin-Sutherland model, and define the QTM and the T -system [21] related to this model. In section 3, we derive traditional TBA equations from the T -system defined in section 2 without using the string hypothesis. In section 4, we derive the NLIE (4.6), (4.7), which are our main results. The normalized fused QTM $\{\mathcal{T}_m^{(a)}(v)\}$ ($a \in \{1, 2, \dots, r\}$; $m \in \mathbb{Z}_{\geq 1}$: the fusion degree of the model) in the Trotter limit $N \rightarrow \infty$ defined in section 2 play the role of the unknown functions of these NLIE. The traditional TBA equations (3.11), (3.13),

(3.16) are integral equations on the variables $\{Y_m^{(a)}(v)\}$, which connects with the ones $\{\mathcal{T}_m^{(a)}(v)\}$ for our new NLIE (4.6) through the relation (3.1) (cf. $\mathcal{T}_m^{(a)}(v) = \lim_{N \rightarrow \infty} \tilde{T}_m^{(a)}(v)$). In this sense, we may view our new NLIE (4.6) as TBA equations on the variables $\{\mathcal{T}_m^{(a)}(v)\}$. For the evaluation of the free energy, we need only the fundamental one $\mathcal{T}_1^{(1)}(v)$. Therefore adopting an infinite number of fused QTM $\{\mathcal{T}_m^{(a)}(v)\}$ as unknown functions seems to be superfluous. In fact we see that these NLIE (4.6) for $m = 1$ form a closed set of equations (4.7), which contains only a *finite* number of unknown functions $\{\mathcal{T}_1^{(a)}(v)\}_{1 \leq a \leq r}$. For $r = 1$, this set of NLIE (4.7) reduces to Takahashi's NLIE [19] for the XXX spin chain. On the other hand, NLIE (4.6) for $m \geq 2$ have never been considered before, and then they are new equations even in the case $r = 1$. Using our new NLIE (4.7), we calculate the high temperature expansion of the free energy in section 5. Section 6 are devoted to concluding remarks. Many calculations in this paper are parallel with the ones for the $osp(1|2s)$ case [12, 22]; but we will describe calculations concisely, not so much for reader's convenience, but for basic importance of the Uimin-Sutherland model.

2 T -system and QTM method

We shall introduce the $sl(r+1)$ Uimin-Sutherland model [24, 25], and define the QTM [5, 6, 7, 8, 9] and the T -system [21] for this model. QTM analyses of the Uimin-Sutherland model can be found in [4, 18]. The classical counterpart of the $sl(r+1)$ Uimin-Sutherland model is a special case of the Perk-Schultz model, whose R -matrix[26] is given as

$$R_{\mu\nu}^{ab}(v) = v\delta_{ab}\delta_{\mu\nu} + \delta_{a\nu}\delta_{\mu b}, \quad (2.1)$$

where $v \in \mathbb{C}$; $a, b, \nu, \mu \in \{1, 2, \dots, r+1\}$. We define the QTM $t_1^{(1)}(v) = ((t_1^{(1)})_{\{\alpha_1, \dots, \alpha_N\}}^{\{\beta_1, \dots, \beta_N\}}(v))$ as

$$(t_1^{(1)})_{\{\alpha_1, \dots, \alpha_N\}}^{\{\beta_1, \dots, \beta_N\}}(v) = \sum_{\{\nu_k\}} \exp\left(\frac{\mu_{\nu_1}}{T}\right) \prod_{k=1}^{\frac{N}{2}} R_{\alpha_{2k-1}, \beta_{2k-1}}^{\nu_{2k-1}, \nu_{2k}}(u+iv) \tilde{R}_{\alpha_{2k}, \beta_{2k}}^{\nu_{2k}, \nu_{2k+1}}(u-iv), \quad (2.2)$$

where $\tilde{R}_{\alpha, \beta}^{\mu, \nu}(v) = R_{\mu, \nu}^{\beta, \alpha}(v)$; N is the Trotter number and assumed to be even; $u = -\frac{J}{TN}$ (T is a temperature; J is a coupling constant, where $J > 0$ (resp. $J < 0$) corresponds to the anti-ferromagnetic (resp. ferromagnetic) regime); $\{\mu_a\}$ are chemical potentials; the Boltzmann constant is set to 1. The free

energy per site is expressed in terms of the largest eigenvalue $T_1^{(1)}(0)$ of the QTM (2.2) at $v = 0$:

$$f = -T \lim_{N \rightarrow \infty} \log T_1^{(1)}(0). \quad (2.3)$$

The eigenvalue formula $T_1^{(1)}(v)$ of the QTM (2.2) is imbedded into the one $T_m^{(a)}(v)$ for a fusion hierarchy of the QTM (cf. Bazhanov-Reshetikhin formula in [29]):

$$T_m^{(a)}(v) = \sum_{\{d_{j,k}\}} \prod_{j=1}^a \prod_{k=1}^m z(d_{j,k}; v - \frac{i}{2}(a - m - 2j + 2k)), \quad (2.4)$$

where the summation is taken over $d_{j,k} \in \{1, 2, \dots, r+1\}$ such that $d_{j,k} \prec d_{j+1,k}$ and $d_{j,k} \preceq d_{j,k+1}$ ($1 \prec 2 \prec \dots \prec r+1$). The functions $\{z(a; v)\}$ are defined as

$$z(a; v) = \psi_a(v) \frac{Q_{a-1}(v - \frac{i}{2}(a+1))Q_a(v - \frac{i}{2}(a-2))}{Q_{a-1}(v - \frac{i}{2}(a-1))Q_a(v - \frac{i}{2}a)} \quad \text{for } a \in \{1, 2, \dots, r+1\}, \quad (2.5)$$

where $Q_a(v) = \prod_{k=1}^{M_a} (v - v_k^{(a)})$; $M_a \in \mathbb{Z}_{\geq 0}$; $Q_0(v) = Q_{r+1}(v) = 1$. The vacuum parts are given as follows

$$\psi_a(v) = e^{\frac{\mu_a}{T}} \phi_+(v + i\delta_{a,r+1}) \phi_-(v - i\delta_{a,1}) \quad \text{for } a \in \{1, 2, \dots, r+1\}, \quad (2.6)$$

where $\phi_{\pm}(v) = (v \pm iu)^{\frac{N}{2}}$. $\{v_k^{(a)}\}$ is a solution of the Bethe ansatz equation (BAE)

$$\frac{\psi_a(v_k^{(a)} + \frac{i}{2}a)}{\psi_{a+1}(v_k^{(a)} + \frac{i}{2}a)} = - \frac{Q_{a-1}(v_k^{(a)} + \frac{i}{2})Q_a(v_k^{(a)} - i)Q_{a+1}(v_k^{(a)} + \frac{i}{2})}{Q_{a-1}(v_k^{(a)} - \frac{i}{2})Q_a(v_k^{(a)} + i)Q_{a+1}(v_k^{(a)} - \frac{i}{2})} \quad (2.7)$$

for $k \in \{1, 2, \dots, M_a\}$ and $a \in \{1, 2, \dots, r\}$.

We can rewrite (2.7) as the Reshetikhin and Wiegmann's BAE [27] in terms of the representation theoretical data.

$$-\prod_{j=1}^N \left(\frac{v_k^{(a)} - \frac{i}{2}b_j^{(a)} - w_j^{(a)}}{v_k^{(a)} + \frac{i}{2}b_j^{(a)} - w_j^{(a)}} \right) = \zeta_a \prod_{b=1}^r \frac{Q_b(v_k^{(a)} - \frac{i}{2}(\alpha_a|\alpha_b))}{Q_b(v_k^{(a)} + \frac{i}{2}(\alpha_a|\alpha_b))} \quad (2.8)$$

for $a \in \{1, 2, \dots, r\}$,

where $(\alpha_a|\alpha_b) = 2\delta_{a,b} - \delta_{a,b+1} - \delta_{a,b-1}$ is the Cartan matrix of $sl(r+1)$; $\{b_j^{(a)}\}$ are the Dynkin labels which characterize the quantum space: $[b_{2j}^{(1)}, b_{2j}^{(2)}, \dots, b_{2j}^{(r)}] =$

$[0, 0, \dots, 1]$, $[b_{2j-1}^{(1)}, b_{2j-1}^{(2)}, \dots, b_{2j-1}^{(r)}] = [1, 0, \dots, 0]$; $\{w_j^{(a)}\}$ are inhomogeneity parameters: $w_{2j-1}^{(a)} = \delta_{a,1} i u$, $w_{2j}^{(a)} = -\delta_{a,r} i(u + \frac{1}{2}g)$; $g = r + 1$ (the dual Coxeter number); $\zeta_a = e^{-\frac{\mu_a - \mu_{a+1}}{T}}$. Note that the Dynkin labels for the even site and the odd site are the ones for conjugate representations each other. The non-negative integers $\{M_a\}$ should satisfy the condition: $\sum_{j=1}^N b_j^{(a)} - \sum_{b=1}^r M_b(\alpha_b | \alpha_a) = \frac{N}{2}(\delta_{a,1} + \delta_{a,r}) + M_{a-1} - 2M_a + M_{a+1} \in \mathbb{Z}_{\geq 0}$, where $M_0 = M_{r+1} = 0$. For the row-to-row transfer matrix, the above parameters take the values: $[b_j^{(1)}, b_j^{(2)}, \dots, b_j^{(r)}] = [1, 0, \dots, 0]$; $w_j^{(a)} = 0$; $\zeta_a = 1$.

For $a \in \{1, 2, \dots, r\}$ and $m \in \mathbb{Z}_{\geq 1}$, we shall normalize (2.4) as $\tilde{T}_m^{(a)}(v) = T_m^{(a)}(v) / \tilde{\mathcal{N}}_m^{(a)}(v)$, where

$$\begin{aligned} \tilde{\mathcal{N}}_m^{(a)}(v) &= \frac{\phi_-(v - \frac{a+m}{2}i)\phi_+(v + \frac{a+m}{2}i)}{\phi_-(v - \frac{a-m}{2}i)\phi_+(v + \frac{a-m}{2}i)} \\ &\times \prod_{j=1}^a \prod_{k=1}^m \phi_-(v - \frac{a-m-2j+2k}{2}i)\phi_+(v - \frac{a-m-2j+2k}{2}i). \end{aligned} \quad (2.9)$$

Note that the poles of $\tilde{T}_m^{(a)}(v)$ from the functions $\{Q_b(v)\}$ (dress part) are spurious under the BAE (2.7). One can show that $\tilde{T}_m^{(a)}(v)$ satisfies the following functional relation.

$$\begin{aligned} \tilde{T}_m^{(a)}(v + \frac{i}{2})\tilde{T}_m^{(a)}(v - \frac{i}{2}) &= \tilde{T}_{m+1}^{(a)}(v)\tilde{T}_{m-1}^{(a)}(v) + \tilde{T}_m^{(a-1)}(v)\tilde{T}_m^{(a+1)}(v), \quad (2.10) \\ \text{for } a &\in \{1, 2, \dots, r\} \quad \text{and} \quad m \in \mathbb{Z}_{\geq 1}, \end{aligned}$$

where

$$\begin{aligned} \tilde{T}_0^{(a)}(v) &= 1 \quad \text{for } a \in \mathbb{Z}_{\geq 1}, \\ \tilde{T}_m^{(0)}(v) &= \frac{\phi_-(v + \frac{m}{2}i)\phi_+(v - \frac{m}{2}i)}{\phi_-(v - \frac{m}{2}i)\phi_+(v + \frac{m}{2}i)} \quad \text{for } m \in \mathbb{Z}_{\geq 1}, \quad (2.11) \\ \tilde{T}_m^{(r+1)}(v) &= e^{\frac{m(\mu_1 + \mu_2 + \dots + \mu_{r+1})}{T}} \quad \text{for } m \in \mathbb{Z}_{\geq 1}. \end{aligned}$$

Except for the vacuum part (the part made from the function (2.6)), this equation has the same form as the $sl(r+1)$ T -system in [21].

3 Traditional TBA equations

In this section, we will transform the T -system (2.10) into the traditional TBA equations by the standard procedure [20, 4]. Similar argument in relation to Stokes multipliers can be found in section 5 in [33].

For $m \in \mathbb{Z}_{\geq 1}$ and $a \in \{1, 2, \dots, r\}$, we define functions:

$$Y_m^{(a)}(v) = \frac{\tilde{T}_{m+1}^{(a)}(v)\tilde{T}_{m-1}^{(a)}(v)}{\tilde{T}_m^{(a-1)}(v)\tilde{T}_m^{(a+1)}(v)} = \frac{T_{m+1}^{(a)}(v)T_{m-1}^{(a)}(v)}{T_m^{(a-1)}(v)T_m^{(a+1)}(v)}. \quad (3.1)$$

By using the T -system (2.10), we can show that (3.1) satisfy the following Y -system:

$$Y_m^{(a)}(v + \frac{i}{2})Y_m^{(a)}(v - \frac{i}{2}) = \frac{(1 + Y_{m+1}^{(a)}(v))(1 + Y_{m-1}^{(a)}(v))}{\prod_{b=1}^r (1 + (Y_m^{(b)}(v))^{-1})^{I_{ab}}} \quad (3.2)$$

for $a \in \{1, 2, \dots, r\}$ and $m \in \mathbb{Z}_{\geq 1}$,

where $I_{ab} = \delta_{a,b+1} + \delta_{a,b-1}$, $Y_0^{(a)}(v) = 0$.

From a numerical analysis for finite N, u, r , we expect that a one-string solution (for every color) in the sector $\frac{N}{2} = M_1 = M_2 = \dots = M_r$ of the BAE (2.7) provides the largest eigenvalue of the QTM (2.2) at $v = 0$ at least for the case: $\mu_1 = \mu_2 = \dots = \mu_{r+1} = 0$ (cf. Figure 1). Hereafter we will consider only this one-string solution. The following conjecture will be valid for this one-string solution (cf. Figures 2, 3).

Conjecture 3.1. *For small u ($|u| \ll 1$), $a \in \{1, 2, \dots, r\}$ and $m \in \mathbb{Z}_{\geq 1}$, every zero $\{\tilde{z}_m^{(a)}\}$ of $\tilde{T}_m^{(a)}(v)$ is located near the lines $\text{Im}v = \pm \frac{m+a}{2}$ at least for the case: $\mu_1 = \mu_2 = \dots = \mu_{r+1} = 0$.*

Based on this conjecture, we will establish the ANZC property (Analytic, NonZero and Constant asymptotics in the limit $|v| \rightarrow \infty$) in some domain for the functions (3.1). We find that $Y_m^{(a)}(v)$ has a constant asymptotics:

$$\lim_{|v| \rightarrow \infty} Y_m^{(a)}(v) = \frac{Q_{m-1}^{(a)}Q_{m+1}^{(a)}}{Q_m^{(a-1)}Q_m^{(a+1)}}, \quad (3.3)$$

where $\{Q_m^{(a)}\}$ is given by (2.4) with a formal setting $Q_a(v) = \phi_{\pm}(v) = 1$. $\{Q_m^{(a)}\}$ is also characterized as a solution of the Q -system [28]:

$$(Q_m^{(a)})^2 = Q_{m-1}^{(a)}Q_{m+1}^{(a)} + Q_m^{(a-1)}Q_m^{(a+1)} \text{ for } a \in \{1, 2, \dots, r\} \text{ and } m \in \mathbb{Z}_{\geq 1}, \quad (3.4)$$

where $Q_0^{(a)} = Q_m^{(0)} = 1$; $Q_m^{(r+1)} = e^{\frac{m(\mu_1 + \mu_2 + \dots + \mu_{r+1})}{T}}$. Note that $Q_m^{(a)}$ coincides with the character of m -th symmetric a -th anti-symmetric tensor representation $V(m\Lambda_a)$ of $gl(r+1)$ if we set $\frac{\mu_a}{T} \rightarrow \epsilon_a$, where $\{\epsilon_a\}$ are orthonormal basis of the dual space of the Cartan subalgebra. In particular, for $\mu_1 = \mu_2 = \dots = \mu_{r+1} = 0$ case, $Q_m^{(a)}$ is given by

$$Q_m^{(a)} = \left(\frac{(m+g)!m!}{(m+a)!(m+g-a)!} \right)^m \prod_{k=1}^m \left\{ \frac{(k+a)(k+g-a)}{k(k+g)} \right\}^k, \quad (3.5)$$




Figure 1: Location of the roots of the BAE for $sl(3)$ case ($N = 12, u = -0.05, \mu_1 = \mu_2 = \mu_3 = 0$), which gives the largest eigenvalue of the QTM $t_1^{(1)}(v)$ at $v = 0$. Both color 1 roots $\{v_k^{(1)}\}$ and color 2 roots $\{v_k^{(2)}\}$ form six one-strings.




Figure 2: Location of zeros of $\tilde{T}_m^{(1)}(v)$ for $sl(3)$ case ($m = 1, 2, 3$, $N = 12, u = -0.05$, $\mu_1 = \mu_2 = \mu_3 = 0$). The zeros recede from the physical strip $\text{Im}v \in [-\frac{1}{2}, \frac{1}{2}]$ as m increases.




Figure 3: Location of zeros of $\tilde{T}_m^{(2)}(v)$ for $sl(3)$ case ($m = 1, 2, 3$, $N = 12, u = -0.05$, $\mu_1 = \mu_2 = \mu_3 = 0$). The zeros recede from the physical strip $\text{Im}v \in [-\frac{1}{2}, \frac{1}{2}]$ as m increases.

and (3.3) has the form

$$\lim_{|v| \rightarrow \infty} Y_m^{(a)}(v) = \frac{m(g+m)}{a(g-a)}. \quad (3.6)$$

(3.5) is the dimension of $V(m\Lambda_a)$. Based on the conjecture 3.1 and (3.3), one can see that the functions $1 + Y_m^{(a)}(v)$, $1 + (Y_m^{(a)}(v))^{-1}$ in the domain $\text{Im} v \in [-\delta, \delta]$ ($0 < \delta \ll 1$) and $Y_m^{(a)}(v)$ for $(a, m) \neq (1, 1)$ in the domain $\text{Im} v \in [-\frac{1}{2}, \frac{1}{2}]$ (physical strip) have the ANZC property at least for the case $\mu_1 = \dots = \mu_{r+1} = 0$; while $Y_1^{(1)}(v)$ has zeros of order $N/2$ at $\pm i(\frac{1}{2} + u)$ if $u < 0$ ($J > 0$), poles of order $N/2$ at $\pm i(\frac{1}{2} - u)$ if $u > 0$ ($J < 0$) in the physical strip. Thus one should modify $Y_m^{(a)}(v)$ as

$$\tilde{Y}_m^{(a)}(v) = Y_m^{(a)}(v) \left\{ \tanh \frac{\pi}{2}(v + i(\frac{1}{2} \pm u)) \tanh \frac{\pi}{2}(v - i(\frac{1}{2} \pm u)) \right\}^{\mp \frac{N\delta_{a,1}\delta_{m,1}}{2}}, \quad (3.7)$$

where the sign $\pm$ is equal to that of $-u$. By using the relation

$$\tanh \frac{\pi}{4}(v + i) \tanh \frac{\pi}{4}(v - i) = 1, \quad (3.8)$$

we shall modify the lhs of the Y -system (3.2) as

$$\tilde{Y}_m^{(a)}(v - \frac{i}{2}) \tilde{Y}_m^{(a)}(v + \frac{i}{2}) = \frac{(1 + Y_{m+1}^{(a)}(v))(1 + Y_{m-1}^{(a)}(v))}{\prod_{b=1}^r (1 + (Y_m^{(b)}(v))^{-1})^{I_{ab}}}, \quad (3.9)$$

for $m \in \mathbb{Z}_{\geq 1}$ and $a \in \{1, 2, \dots, r\}$.

This modified Y -system has the ANZC property. Then one can transform (3.9) into nonlinear integral equations by a standard procedure.

$$\begin{aligned} \log Y_m^{(a)}(v) &= \pm \frac{N\delta_{a,1}\delta_{m,1}}{2} \log \left\{ \tanh \frac{\pi}{2}(v + i(\frac{1}{2} \pm u)) \tanh \frac{\pi}{2}(v - i(\frac{1}{2} \pm u)) \right\} \\ &+ K * \log \left\{ \frac{(1 + Y_{m-1}^{(a)})(1 + Y_{m+1}^{(a)})}{\prod_{b=1}^r (1 + (Y_m^{(b)})^{-1})^{I_{ab}}} \right\} (v) \end{aligned} \quad (3.10)$$

for $a \in \{1, 2, \dots, r\}$ and $m \in \mathbb{Z}_{\geq 1}$,

where $Y_0^{(a)}(v) = 0$ and $*$ is the convolution. After the Trotter limit $N \rightarrow \infty$ with $u = -\frac{J}{NT}$, we obtain a TBA equation

$$\begin{aligned} \log Y_m^{(a)}(v) &= -\frac{\pi J \delta_{a,1} \delta_{m,1}}{T \cosh \pi v} + K * \log \left\{ \frac{(1 + Y_{m-1}^{(a)})(1 + Y_{m+1}^{(a)})}{\prod_{b=1}^r (1 + (Y_m^{(b)})^{-1})^{I_{ab}}} \right\} (v) \end{aligned} \quad (3.11)$$

for $a \in \{1, 2, \dots, r\}$ and $m \in \mathbb{Z}_{\geq 1}$,

where $Y_0^{(a)}(v) := 0$ and the kernel is

$$K(v) = \frac{1}{2 \cosh \pi v}. \quad (3.12)$$

One can also rewrite (3.11) as

$$\begin{aligned} -\frac{\pi J \delta_{a1} \delta_{m1}}{T \cosh \pi v} &= \sum_{n=1}^{\infty} K^{m,n} * \log(1 + Y_n^{(a)})(v) \\ &\quad - \sum_{b=1}^r \sum_{n=1}^{\infty} J_{a,b}^{m,n} * \log(1 + (Y_n^{(b)})^{-1})(v), \end{aligned} \quad (3.13)$$

where

$$\begin{aligned} K^{m,n}(v) &= \delta_{m,n} \delta(v) - (\delta_{m,n-1} + \delta_{m,n+1}) K(v), \\ J_{a,b}^{n,k}(v) &= (\delta_{a,b} \delta(v) - (\delta_{a,b-1} + \delta_{a,b+1}) K(v)) \delta_{n,k}. \end{aligned} \quad (3.14)$$

We find that the TBA equation (3.13) coincides with the one from the string hypothesis in [29] after suitable modification. By using the following relations

$$\begin{aligned} A^{a,d}(v) &= \sum_{l=1}^{\min(a,d)} G_{|a-d|+2l-1}(v), \\ G_a(v) &= \frac{1}{g} \frac{\sin \frac{(g-a)\pi}{g}}{\cos \frac{(g-a)\pi}{g} + \cosh \frac{2\pi v}{g}}, \\ \hat{A}^{a,d}(k) &= \int_{-\infty}^{\infty} dv A^{a,d}(v) e^{-ikv} = \frac{\sinh \left(\frac{\min(a,d)}{2} k \right) \sinh \left(\frac{g-\min(a,d)}{2} k \right)}{\sinh \left(\frac{k}{2} \right) \sinh \left(\frac{g}{2} k \right)}, \\ \sum_{l=1}^r \hat{A}^{m,l}(k) \hat{K}^{l,n}(k) &= \frac{\delta_{m,n}}{2 \cosh \left(\frac{k}{2} \right)}, \\ \hat{K}^{m,n}(k) &= \delta_{m,n} - \frac{\delta_{m,n-1} + \delta_{m,n+1}}{2 \cosh \frac{k}{2}}, \end{aligned} \quad (3.15)$$

we can also rewrite the TBA equation (3.11) as

$$\begin{aligned} \log Y_m^{(a)}(v) &= -\frac{2\pi J \delta_{m,1}}{T} G_a(v) \\ &\quad + \sum_{b=1}^r A^{a,b} * \log \left\{ \frac{(1 + Y_{m-1}^{(b)})(1 + Y_{m+1}^{(b)})}{\prod_{d=1}^r (1 + Y_m^{(d)})^{I_{bd}}} \right\} (v) \\ &\quad \text{for } a \in \{1, 2, \dots, r\} \text{ and } m \in \mathbb{Z}_{\geq 1}, \end{aligned} \quad (3.16)$$

where $Y_0^{(a)}(v) = 0$. This TBA equation (3.16) does not contain $(Y_m^{(a)})^{-1}$, which contrasts with (3.11), (3.13). We can derive the following relation from (2.10), (3.1) and (3.15).

$$\begin{aligned} \log \tilde{T}_1^{(1)}(v) &= \sum_{b=1}^r G_b * \log(1 + Y_1^{(b)})(v) \\ &\quad - 2N \int_0^\infty dk \frac{e^{-\frac{1}{2}k} \sinh(ku) \cos(kv) \sinh(\frac{g-1}{2}k)}{k \sinh(\frac{g}{2}k)}. \end{aligned} \quad (3.17)$$

After the Trotter limit $N \rightarrow \infty$ with $u = -\frac{J}{NT}$, we have the free energy per site.

$$\begin{aligned} f &= -T \lim_{N \rightarrow \infty} \log T_1^{(1)}(0) = J - T \lim_{N \rightarrow \infty} \log \tilde{T}_1^{(1)}(0) \\ &= J \left\{ \frac{2}{g} \left(\gamma + \psi\left(\frac{1}{g}\right) \right) + 1 \right\} - T \sum_{a=1}^r \int_{-\infty}^\infty dv G_a(v) \log(1 + Y_1^{(a)}(v)), \end{aligned} \quad (3.18)$$

where $\psi(z)$ is the digamma function

$$\psi(z) = \frac{d}{dz} \log \Gamma(z), \quad \gamma = -\psi(1). \quad (3.19)$$

4 New nonlinear integral equations

It was pointed out [13] that Takahashi's NLIE for the XXZ -model [19] can be rederived from the T -system of the QTM. In this section, we shall derive our new NLIE (4.6) and (4.7) from the T -system (2.10).

$\tilde{T}_m^{(a)}(v)$ has poles only at $\pm \tilde{\beta}_m^{(a)}$: $\tilde{\beta}_m^{(a)} = \frac{m+a}{2}i + iu$. These poles are of order $N/2$ at most. In addition, $\lim_{|v| \rightarrow \infty} \tilde{T}_m^{(a)}(v) = Q_m^{(a)}$ is a finite number (cf. (3.5)). Thus we must put

$$\tilde{T}_m^{(a)}(v) = Q_m^{(a)} + \sum_{j=1}^{\frac{N}{2}} \left\{ \frac{b_{m,j}^{(a)}}{(v - \tilde{\beta}_m^{(a)})^j} + \frac{\bar{b}_{m,j}^{(a)}}{(v + \tilde{\beta}_m^{(a)})^j} \right\}, \quad (4.1)$$

where the coefficients $b_{m,j}^{(a)}, \bar{b}_{m,j}^{(a)} \in \mathbb{C}$ are given as follows.

$$\begin{aligned} b_{m,j}^{(a)} &= \oint_{C_m^{(a)}} \frac{dv}{2\pi i} \tilde{T}_m^{(a)}(v) (v - \tilde{\beta}_m^{(a)})^{j-1}, \\ \bar{b}_{m,j}^{(a)} &= \oint_{\bar{C}_m^{(a)}} \frac{dv}{2\pi i} \tilde{T}_m^{(a)}(v) (v + \tilde{\beta}_m^{(a)})^{j-1}, \end{aligned} \quad (4.2)$$

where the contour $C_m^{(a)}$ is a counterclockwise closed loop around $\tilde{\beta}_m^{(a)}$ which does not surround $-\tilde{\beta}_m^{(a)}$; the contour $\overline{C}_m^{(a)}$ is a counterclockwise closed loop around $-\tilde{\beta}_m^{(a)}$ which does not surround $\tilde{\beta}_m^{(a)}$. Using the T -system (2.10), one can modify (4.2) as

$$\begin{aligned} b_{m,j}^{(a)} &= \oint_{C_m^{(a)}} \frac{dv}{2\pi i} \left\{ \frac{\tilde{T}_{m-1}^{(a)}(v - \frac{i}{2})\tilde{T}_{m+1}^{(a)}(v - \frac{i}{2})}{\tilde{T}_m^{(a)}(v - i)} \right. \\ &\quad \left. + \frac{\tilde{T}_m^{(a-1)}(v - \frac{i}{2})\tilde{T}_m^{(a+1)}(v - \frac{i}{2})}{\tilde{T}_m^{(a)}(v - i)} \right\} (v - \tilde{\beta}_m^{(a)})^{j-1}, \\ \bar{b}_{m,j}^{(a)} &= \oint_{\overline{C}_m^{(a)}} \frac{dv}{2\pi i} \left\{ \frac{\tilde{T}_{m-1}^{(a)}(v + \frac{i}{2})\tilde{T}_{m+1}^{(a)}(v + \frac{i}{2})}{\tilde{T}_m^{(a)}(v + i)} \right. \\ &\quad \left. + \frac{\tilde{T}_m^{(a-1)}(v + \frac{i}{2})\tilde{T}_m^{(a+1)}(v + \frac{i}{2})}{\tilde{T}_m^{(a)}(v + i)} \right\} (v + \tilde{\beta}_m^{(a)})^{j-1}. \quad (4.3) \end{aligned}$$

Substituting (4.3) into (4.1), we obtain

$$\begin{aligned} \tilde{T}_m^{(a)}(v) &= Q_m^{(a)} \\ &+ \oint_{C_m^{(a)}} \frac{dy}{2\pi i} \frac{1 - \left(\frac{y}{v - \tilde{\beta}_m^{(a)}}\right)^{\frac{N}{2}}}{v - y - \tilde{\beta}_m^{(a)}} \left\{ \frac{\tilde{T}_{m-1}^{(a)}(y + \tilde{\beta}_m^{(a)} - \frac{i}{2})\tilde{T}_{m+1}^{(a)}(y + \tilde{\beta}_m^{(a)} - \frac{i}{2})}{\tilde{T}_m^{(a)}(y + \tilde{\beta}_m^{(a)} - i)} \right. \\ &\quad \left. + \frac{\tilde{T}_m^{(a-1)}(y + \tilde{\beta}_m^{(a)} - \frac{i}{2})\tilde{T}_m^{(a+1)}(y + \tilde{\beta}_m^{(a)} - \frac{i}{2})}{\tilde{T}_m^{(a)}(y + \tilde{\beta}_m^{(a)} - i)} \right\} \\ &+ \oint_{\overline{C}_m^{(a)}} \frac{dy}{2\pi i} \frac{1 - \left(\frac{y}{v + \tilde{\beta}_m^{(a)}}\right)^{\frac{N}{2}}}{v - y + \tilde{\beta}_m^{(a)}} \left\{ \frac{\tilde{T}_{m-1}^{(a)}(y - \tilde{\beta}_m^{(a)} + \frac{i}{2})\tilde{T}_{m+1}^{(a)}(y - \tilde{\beta}_m^{(a)} + \frac{i}{2})}{\tilde{T}_m^{(a)}(y - \tilde{\beta}_m^{(a)} + i)} \right. \\ &\quad \left. + \frac{\tilde{T}_m^{(a-1)}(y - \tilde{\beta}_m^{(a)} + \frac{i}{2})\tilde{T}_m^{(a+1)}(y - \tilde{\beta}_m^{(a)} + \frac{i}{2})}{\tilde{T}_m^{(a)}(y - \tilde{\beta}_m^{(a)} + i)} \right\}, \\ &\text{for } a \in \{1, 2, \dots, r\} \text{ and } m \in \mathbb{Z}_{\geq 1}, \quad (4.4) \end{aligned}$$

where the contour $C_m^{(a)}$ (resp. $\overline{C}_m^{(a)}$) is a counterclockwise closed loop around 0 which does not surround $-2\tilde{\beta}_m^{(a)}$ (resp. $2\tilde{\beta}_m^{(a)}$). The first term and the second term in the first bracket $\{\dots\}$ in (4.4) have a common singularity at 0; while for $m = 1$, this singularity from the first term disappears since $\tilde{T}_0^{(a)}(y) = 1$. Thus for $m = 1$, the contribution to the contour integral from the first term in the first bracket $\{\dots\}$ in (4.4) vanishes if the contour $C_1^{(a)}$ does not surround the singularity at $\tilde{z}_1^{(a)} - \tilde{\beta}_1^{(a)} + i$ (cf. Conjecture 3.1). This

is also the case with the second bracket $\{\cdots\}$ in (4.4). Therefore for $m = 1$, (4.4) reduces to

$$\begin{aligned}
\tilde{T}_1^{(a)}(v) &= Q_1^{(a)} \\
&+ \oint_{C_1^{(a)}} \frac{dy}{2\pi i} \frac{1 - \left(\frac{y}{v - \tilde{\beta}_1^{(a)}}\right)^{\frac{N}{2}}}{v - y - \tilde{\beta}_1^{(a)}} \frac{\tilde{T}_1^{(a-1)}(y + \tilde{\beta}_1^{(a)} - \frac{i}{2}) \tilde{T}_1^{(a+1)}(y + \tilde{\beta}_1^{(a)} - \frac{i}{2})}{\tilde{T}_1^{(a)}(y + \tilde{\beta}_1^{(a)} - i)} \\
&+ \oint_{\bar{C}_1^{(a)}} \frac{dy}{2\pi i} \frac{1 - \left(\frac{y}{v + \tilde{\beta}_1^{(a)}}\right)^{\frac{N}{2}}}{v - y + \tilde{\beta}_1^{(a)}} \frac{\tilde{T}_1^{(a-1)}(y - \tilde{\beta}_1^{(a)} + \frac{i}{2}) \tilde{T}_1^{(a+1)}(y - \tilde{\beta}_1^{(a)} + \frac{i}{2})}{\tilde{T}_1^{(a)}(y - \tilde{\beta}_1^{(a)} + i)} \\
&\quad \text{for } a \in \{1, 2, \dots, r\}, \tag{4.5}
\end{aligned}$$

where the contour $C_1^{(a)}$ (resp. $\bar{C}_1^{(a)}$) is a counterclockwise closed loop around 0 which does not surround $\tilde{z}_1^{(a)} - \tilde{\beta}_1^{(a)} + i$, $-2\tilde{\beta}_1^{(a)}$ (resp. $\tilde{z}_1^{(a)} + \tilde{\beta}_1^{(a)} - i$, $2\tilde{\beta}_1^{(a)}$). Now we shall take the Trotter limit $N \rightarrow \infty$ in (4.4).

$$\begin{aligned}
\mathcal{T}_m^{(a)}(v) &= Q_m^{(a)} \\
&+ \oint_{C_m^{(a)}} \frac{dy}{2\pi i} \frac{1}{v - y - \beta_m^{(a)}} \left\{ \frac{\mathcal{T}_{m-1}^{(a)}(y + \beta_m^{(a)} - \frac{i}{2}) \mathcal{T}_{m+1}^{(a)}(y + \beta_m^{(a)} - \frac{i}{2})}{\mathcal{T}_m^{(a)}(y + \beta_m^{(a)} - i)} \right. \\
&\quad \left. + \frac{\mathcal{T}_m^{(a-1)}(y + \beta_m^{(a)} - \frac{i}{2}) \mathcal{T}_m^{(a+1)}(y + \beta_m^{(a)} - \frac{i}{2})}{\mathcal{T}_m^{(a)}(y + \beta_m^{(a)} - i)} \right\} \\
&+ \oint_{\bar{C}_m^{(a)}} \frac{dy}{2\pi i} \frac{1}{v - y + \beta_m^{(a)}} \left\{ \frac{\mathcal{T}_{m-1}^{(a)}(y - \beta_m^{(a)} + \frac{i}{2}) \mathcal{T}_{m+1}^{(a)}(y - \beta_m^{(a)} + \frac{i}{2})}{\mathcal{T}_m^{(a)}(y - \beta_m^{(a)} + i)} \right. \\
&\quad \left. + \frac{\mathcal{T}_m^{(a-1)}(y - \beta_m^{(a)} + \frac{i}{2}) \mathcal{T}_m^{(a+1)}(y - \beta_m^{(a)} + \frac{i}{2})}{\mathcal{T}_m^{(a)}(y - \beta_m^{(a)} + i)} \right\} \\
&\quad \text{for } a \in \{1, 2, \dots, r\} \text{ and } m \in \mathbb{Z}_{\geq 1}, \tag{4.6}
\end{aligned}$$

where $\mathcal{T}_m^{(a)}(v) := \lim_{N \rightarrow \infty} \tilde{T}_m^{(a)}(v)$; $\beta_m^{(a)} := \lim_{N \rightarrow \infty} \tilde{\beta}_m^{(a)} = \frac{m+a}{2}i$; $\mathcal{T}_m^{(r+1)}(v) = e^{\frac{m(\mu_1 + \dots + \mu_{r+1})}{T}}$; $\mathcal{T}_0^{(a)}(v) = 1$; $\mathcal{T}_m^{(0)}(v) = \exp\left(\frac{mJ}{(v^2 + \frac{m^2}{4})T}\right)$; the contour $C_m^{(a)}$ (resp. $\bar{C}_m^{(a)}$) is a counterclockwise closed loop around 0 which satisfies the condition $|y| < |v - \beta_m^{(a)}|$ (resp. $|y| < |v + \beta_m^{(a)}|$) and does not surround $-2\beta_m^{(a)}$ (resp. $2\beta_m^{(a)}$). In particular for $m = 1$, we obtain a system of NLIE, which contains

only a *finite* number of unknown functions $\{\mathcal{T}_1^{(a)}(v)\}_{1 \leq a \leq r}$:

$$\begin{aligned} \mathcal{T}_1^{(a)}(v) = Q_1^{(a)} &+ \oint_{C_1^{(a)}} \frac{dy}{2\pi i} \frac{\mathcal{T}_1^{(a-1)}(y + \beta_1^{(a)} - \frac{i}{2}) \mathcal{T}_1^{(a+1)}(y + \beta_1^{(a)} - \frac{i}{2})}{(v - y - \beta_1^{(a)}) \mathcal{T}_1^{(a)}(y + \beta_1^{(a)} - i)} \\ &+ \oint_{\bar{C}_1^{(a)}} \frac{dy}{2\pi i} \frac{\mathcal{T}_1^{(a-1)}(y - \beta_1^{(a)} + \frac{i}{2}) \mathcal{T}_1^{(a+1)}(y - \beta_1^{(a)} + \frac{i}{2})}{(v - y + \beta_1^{(a)}) \mathcal{T}_1^{(a)}(y - \beta_1^{(a)} + i)} \\ &\text{for } a \in \{1, 2, \dots, r\}, \end{aligned} \quad (4.7)$$

where $\mathcal{T}_1^{(r+1)}(v) = e^{\frac{\mu_1 + \dots + \mu_{r+1}}{T}}$; the contour $C_1^{(a)}$ (resp. $\bar{C}_1^{(a)}$) is a counter-clockwise closed loop around 0 which satisfies the condition $|y| < |v - \beta_1^{(a)}|$ (resp. $|y| < |v + \beta_1^{(a)}|$) and does not surround $z_1^{(a)} - \beta_1^{(a)} + i$, $-2\beta_1^{(a)}$ (resp. $z_1^{(a)} + \beta_1^{(a)} - i$, $2\beta_1^{(a)}$); $z_1^{(a)} = \lim_{N \rightarrow \infty} \tilde{z}_1^{(a)}$. One can calculate the free energy per site f by using (4.7) and the relation

$$f = J - T \log \mathcal{T}_1^{(1)}(0). \quad (4.8)$$

We need not use (4.6) (for $m \in \mathbb{Z}_{\geq 2}$) to obtain the free energy per site. In addition, $\{\mathcal{T}_m^{(a)}(v)\}_{1 \leq a \leq r; m \in \mathbb{Z}_{\geq 1}}$ is given in terms of the solution of (4.7) $\{\mathcal{T}_1^{(a)}(v)\}_{1 \leq a \leq r}$ based on a Jacobi-Trudi formula [29]. However, (4.6) has theoretical significance since it manifests the relation between our new NLIE and the traditional TBA equations (3.11), (3.13), (3.16). In fact, (4.6) and (4.7) are related to the traditional TBA equations (3.11), (3.13), (3.16) through the relation (3.1) in the Trotter limit.

5 High temperature expansion

In this section, we shall calculate the high temperature expansion of the free energy (4.8) for $r = 1, 2, 3$ by our new NLIE (4.7). As for the XXX -model case, the high temperature expansion of the free energy is calculated [30] by Takahashi's NLIE up to order 100. We assume the following expansion for large T :

$$\mathcal{T}_1^{(a)}(v) = \exp \left(\sum_{n=0}^{\infty} b_n^{(a)}(v) \left(\frac{J}{T} \right)^n \right). \quad (5.1)$$

In contrasts with the XXX -model case [30], we need further assumption

$$b_n^{(a)}(v) = \sum_{j=0}^{n-1} \frac{c_{n,j}^{(a)} v^{2j}}{(v^2 + \frac{(a+1)^2}{4})^n}, \quad (5.2)$$

where $c_{n,j}^{(a)} \in \mathbb{C}$. Substituting (5.1) into (4.7), we can obtain the coefficients $\{b_n^{(a)}(v)\}$. Here we just enumerate first few of them. $b_0^{(a)}(v)$ has the form $b_0^{(a)}(v) = \log Q_1^{(a)}$. Other coefficients are given as follows.
 $sl(2)$ case:

$$\begin{aligned}
c_{1,0}^{(1)} &= \frac{2 Q_1^{(2)}}{Q_1^{(1)2}}, \\
c_{2,0}^{(1)} &= \frac{3 Q_1^{(2)}}{Q_1^{(1)2}} - \frac{6 Q_1^{(2)2}}{Q_1^{(1)4}}, \\
c_{2,1}^{(1)} &= \frac{Q_1^{(2)}}{Q_1^{(1)2}} - \frac{4 Q_1^{(2)2}}{Q_1^{(1)4}}, \\
c_{3,0}^{(1)} &= \frac{10 Q_1^{(2)}}{3 Q_1^{(1)2}} - \frac{18 Q_1^{(2)2}}{Q_1^{(1)4}} + \frac{80 Q_1^{(2)3}}{3 Q_1^{(1)6}}, \\
c_{3,1}^{(1)} &= \frac{3 Q_1^{(2)}}{Q_1^{(1)2}} - \frac{22 Q_1^{(2)2}}{Q_1^{(1)4}} + \frac{40 Q_1^{(2)3}}{Q_1^{(1)6}}, \\
c_{3,2}^{(1)} &= \frac{Q_1^{(2)}}{Q_1^{(1)2}} - \frac{8 Q_1^{(2)2}}{Q_1^{(1)4}} + \frac{16 Q_1^{(2)3}}{Q_1^{(1)6}}.
\end{aligned} \tag{5.3}$$

$sl(3)$ case:

$$\begin{aligned}
c_{1,0}^{(1)} &= \frac{2 Q_1^{(2)}}{Q_1^{(1)2}}, \\
c_{2,0}^{(1)} &= \frac{3 Q_1^{(2)}}{Q_1^{(1)2}} - \frac{6 Q_1^{(2)2}}{Q_1^{(1)4}} + \frac{3 Q_1^{(3)}}{Q_1^{(1)3}}, \\
c_{2,1}^{(1)} &= \frac{Q_1^{(2)}}{Q_1^{(1)2}} - \frac{4 Q_1^{(2)2}}{Q_1^{(1)4}} + \frac{3 Q_1^{(3)}}{Q_1^{(1)3}}, \\
c_{3,0}^{(1)} &= \frac{10 Q_1^{(2)}}{3 Q_1^{(1)2}} - \frac{18 Q_1^{(2)2}}{Q_1^{(1)4}} + \frac{80 Q_1^{(2)3}}{3 Q_1^{(1)6}} + \frac{8 Q_1^{(3)}}{Q_1^{(1)3}} - \frac{24 Q_1^{(2)} Q_1^{(3)}}{Q_1^{(1)5}}, \\
c_{3,1}^{(1)} &= \frac{3 Q_1^{(2)}}{Q_1^{(1)2}} - \frac{22 Q_1^{(2)2}}{Q_1^{(1)4}} + \frac{40 Q_1^{(2)3}}{Q_1^{(1)6}} + \frac{13 Q_1^{(3)}}{Q_1^{(1)3}} - \frac{42 Q_1^{(2)} Q_1^{(3)}}{Q_1^{(1)5}}, \\
c_{3,2}^{(1)} &= \frac{Q_1^{(2)}}{Q_1^{(1)2}} - \frac{8 Q_1^{(2)2}}{Q_1^{(1)4}} + \frac{16 Q_1^{(2)3}}{Q_1^{(1)6}} + \frac{5 Q_1^{(3)}}{Q_1^{(1)3}} - \frac{18 Q_1^{(2)} Q_1^{(3)}}{Q_1^{(1)5}},
\end{aligned} \tag{5.4}$$

$$\begin{aligned}
c_{1,0}^{(2)} &= \frac{3 Q_1^{(3)}}{Q_1^{(1)} Q_1^{(2)}}, \\
c_{2,0}^{(2)} &= \frac{-27 Q_1^{(3)}}{2 Q_1^{(1)3}} + \frac{9 Q_1^{(3)}}{Q_1^{(1)} Q_1^{(2)}} - \frac{9 Q_1^{(3)2}}{2 Q_1^{(1)2} Q_1^{(2)2}}, \\
c_{2,1}^{(2)} &= \frac{-6 Q_1^{(3)}}{Q_1^{(1)3}} + \frac{2 Q_1^{(3)}}{Q_1^{(1)} Q_1^{(2)}}, \\
c_{3,0}^{(2)} &= \frac{-1377 Q_1^{(3)}}{16 Q_1^{(1)3}} + \frac{171 Q_1^{(3)}}{8 Q_1^{(1)} Q_1^{(2)}} + \frac{243 Q_1^{(2)} Q_1^{(3)}}{2 Q_1^{(1)5}} - \frac{27 Q_1^{(3)2}}{Q_1^{(1)2} Q_1^{(2)2}} \\
&\quad - \frac{81 Q_1^{(3)2}}{16 Q_1^{(1)4} Q_1^{(2)}} + \frac{9 Q_1^{(3)3}}{Q_1^{(1)3} Q_1^{(2)3}}, \\
c_{3,1}^{(2)} &= \frac{-135 Q_1^{(3)}}{2 Q_1^{(1)3}} + \frac{12 Q_1^{(3)}}{Q_1^{(1)} Q_1^{(2)}} + \frac{108 Q_1^{(2)} Q_1^{(3)}}{Q_1^{(1)5}} - \frac{6 Q_1^{(3)2}}{Q_1^{(1)2} Q_1^{(2)2}} - \frac{45 Q_1^{(3)2}}{2 Q_1^{(1)4} Q_1^{(2)}}, \\
c_{3,2}^{(2)} &= \frac{-13 Q_1^{(3)}}{Q_1^{(1)3}} + \frac{2 Q_1^{(3)}}{Q_1^{(1)} Q_1^{(2)}} + \frac{24 Q_1^{(2)} Q_1^{(3)}}{Q_1^{(1)5}} - \frac{9 Q_1^{(3)2}}{Q_1^{(1)4} Q_1^{(2)}}.
\end{aligned} \tag{5.5}$$

$sl(4)$ case:

$$\begin{aligned}
c_{1,0}^{(1)} &= \frac{2 Q_1^{(2)}}{Q_1^{(1)2}}, \\
c_{2,0}^{(1)} &= \frac{3 Q_1^{(2)}}{Q_1^{(1)2}} - \frac{6 Q_1^{(2)2}}{Q_1^{(1)4}} + \frac{3 Q_1^{(3)}}{Q_1^{(1)3}}, \\
c_{2,1}^{(1)} &= \frac{Q_1^{(2)}}{Q_1^{(1)2}} - \frac{4 Q_1^{(2)2}}{Q_1^{(1)4}} + \frac{3 Q_1^{(3)}}{Q_1^{(1)3}}, \\
c_{3,0}^{(1)} &= \frac{10 Q_1^{(2)}}{3 Q_1^{(1)2}} - \frac{18 Q_1^{(2)2}}{Q_1^{(1)4}} + \frac{80 Q_1^{(2)3}}{3 Q_1^{(1)6}} + \frac{8 Q_1^{(3)}}{Q_1^{(1)3}} - \frac{24 Q_1^{(2)} Q_1^{(3)}}{Q_1^{(1)5}} + \frac{4 Q_1^{(4)}}{Q_1^{(1)4}}, \\
c_{3,1}^{(1)} &= \frac{3 Q_1^{(2)}}{Q_1^{(1)2}} - \frac{22 Q_1^{(2)2}}{Q_1^{(1)4}} + \frac{40 Q_1^{(2)3}}{Q_1^{(1)6}} + \frac{13 Q_1^{(3)}}{Q_1^{(1)3}} - \frac{42 Q_1^{(2)} Q_1^{(3)}}{Q_1^{(1)5}} + \frac{8 Q_1^{(4)}}{Q_1^{(1)4}}, \\
c_{3,2}^{(1)} &= \frac{Q_1^{(2)}}{Q_1^{(1)2}} - \frac{8 Q_1^{(2)2}}{Q_1^{(1)4}} + \frac{16 Q_1^{(2)3}}{Q_1^{(1)6}} + \frac{5 Q_1^{(3)}}{Q_1^{(1)3}} - \frac{18 Q_1^{(2)} Q_1^{(3)}}{Q_1^{(1)5}} + \frac{4 Q_1^{(4)}}{Q_1^{(1)4}},
\end{aligned} \tag{5.6}$$

$$\begin{aligned}
c_{1,0}^{(2)} &= \frac{3 Q_1^{(3)}}{Q_1^{(1)} Q_1^{(2)}}, \\
c_{2,0}^{(2)} &= \frac{-27 Q_1^{(3)}}{2 Q_1^{(1)3}} + \frac{9 Q_1^{(3)}}{Q_1^{(1)} Q_1^{(2)}} - \frac{9 Q_1^{(3)2}}{2 Q_1^{(1)2} Q_1^{(2)2}} + \frac{9 Q_1^{(4)}}{Q_1^{(1)2} Q_1^{(2)}}, \\
c_{2,1}^{(2)} &= \frac{-6 Q_1^{(3)}}{Q_1^{(1)3}} + \frac{2 Q_1^{(3)}}{Q_1^{(1)} Q_1^{(2)}} + \frac{4 Q_1^{(4)}}{Q_1^{(1)2} Q_1^{(2)}}, \\
c_{3,0}^{(2)} &= \frac{-1377 Q_1^{(3)}}{16 Q_1^{(1)3}} + \frac{171 Q_1^{(3)}}{8 Q_1^{(1)} Q_1^{(2)}} + \frac{243 Q_1^{(2)} Q_1^{(3)}}{2 Q_1^{(1)5}} - \frac{27 Q_1^{(3)2}}{Q_1^{(1)2} Q_1^{(2)2}} - \frac{81 Q_1^{(3)2}}{16 Q_1^{(1)4} Q_1^{(2)}} \\
&\quad + \frac{9 Q_1^{(3)3}}{Q_1^{(1)3} Q_1^{(2)3}} - \frac{81 Q_1^{(4)}}{Q_1^{(1)4}} + \frac{783 Q_1^{(4)}}{16 Q_1^{(1)2} Q_1^{(2)}} - \frac{27 Q_1^{(3)} Q_1^{(4)}}{Q_1^{(1)3} Q_1^{(2)2}}, \\
c_{3,1}^{(2)} &= \frac{-135 Q_1^{(3)}}{2 Q_1^{(1)3}} + \frac{12 Q_1^{(3)}}{Q_1^{(1)} Q_1^{(2)}} + \frac{108 Q_1^{(2)} Q_1^{(3)}}{Q_1^{(1)5}} - \frac{6 Q_1^{(3)2}}{Q_1^{(1)2} Q_1^{(2)2}} \\
&\quad - \frac{45 Q_1^{(3)2}}{2 Q_1^{(1)4} Q_1^{(2)}} - \frac{72 Q_1^{(4)}}{Q_1^{(1)4}} + \frac{75 Q_1^{(4)}}{2 Q_1^{(1)2} Q_1^{(2)}} - \frac{12 Q_1^{(3)} Q_1^{(4)}}{Q_1^{(1)3} Q_1^{(2)2}}, \\
c_{3,2}^{(2)} &= \frac{-13 Q_1^{(3)}}{Q_1^{(1)3}} + \frac{2 Q_1^{(3)}}{Q_1^{(1)} Q_1^{(2)}} + \frac{24 Q_1^{(2)} Q_1^{(3)}}{Q_1^{(1)5}} - \frac{9 Q_1^{(3)2}}{Q_1^{(1)4} Q_1^{(2)}} - \frac{16 Q_1^{(4)}}{Q_1^{(1)4}} + \frac{7 Q_1^{(4)}}{Q_1^{(1)2} Q_1^{(2)}}, \\
&\hspace{15em} (5.7)
\end{aligned}$$

$$\begin{aligned}
c_{1,0}^{(3)} &= \frac{4 Q_1^{(4)}}{Q_1^{(1)} Q_1^{(3)}}, \\
c_{2,0}^{(3)} &= \frac{20 Q_1^{(4)}}{Q_1^{(1)} Q_1^{(3)}} - \frac{32 Q_1^{(2)} Q_1^{(4)}}{Q_1^{(1)3} Q_1^{(3)}} - \frac{8 Q_1^{(4)2}}{Q_1^{(1)2} Q_1^{(3)2}}, \\
c_{2,1}^{(3)} &= \frac{3 Q_1^{(4)}}{Q_1^{(1)} Q_1^{(3)}} - \frac{8 Q_1^{(2)} Q_1^{(4)}}{Q_1^{(1)3} Q_1^{(3)}}, \\
c_{3,0}^{(3)} &= \frac{-192 Q_1^{(4)}}{Q_1^{(1)4}} + \frac{248 Q_1^{(4)}}{3 Q_1^{(1)} Q_1^{(3)}} - \frac{352 Q_1^{(2)} Q_1^{(4)}}{Q_1^{(1)3} Q_1^{(3)}} + \frac{512 Q_1^{(2)2} Q_1^{(4)}}{Q_1^{(1)5} Q_1^{(3)}} \\
&\quad - \frac{80 Q_1^{(4)2}}{Q_1^{(1)2} Q_1^{(3)2}} + \frac{128 Q_1^{(2)} Q_1^{(4)2}}{Q_1^{(1)4} Q_1^{(3)2}} + \frac{64 Q_1^{(4)3}}{3 Q_1^{(1)3} Q_1^{(3)3}}, \\
c_{3,1}^{(3)} &= \frac{-96 Q_1^{(4)}}{Q_1^{(1)4}} + \frac{30 Q_1^{(4)}}{Q_1^{(1)} Q_1^{(3)}} - \frac{160 Q_1^{(2)} Q_1^{(4)}}{Q_1^{(1)3} Q_1^{(3)}} + \frac{256 Q_1^{(2)2} Q_1^{(4)}}{Q_1^{(1)5} Q_1^{(3)}} \\
&\quad - \frac{12 Q_1^{(4)2}}{Q_1^{(1)2} Q_1^{(3)2}} + \frac{32 Q_1^{(2)} Q_1^{(4)2}}{Q_1^{(1)4} Q_1^{(3)2}}, \\
c_{3,2}^{(3)} &= \frac{-12 Q_1^{(4)}}{Q_1^{(1)4}} + \frac{3 Q_1^{(4)}}{Q_1^{(1)} Q_1^{(3)}} - \frac{18 Q_1^{(2)} Q_1^{(4)}}{Q_1^{(1)3} Q_1^{(3)}} + \frac{32 Q_1^{(2)2} Q_1^{(4)}}{Q_1^{(1)5} Q_1^{(3)}}.
\end{aligned} \tag{5.8}$$

When $Q_1^{(a)}$ are suitably chosen, (5.3) recover the known results[30]. One see that these coefficients are expressed in terms of the solutions of the Q -system (3.4). And thus a solution of the T -system is given in terms of solutions of the Q -system. The T -system is a Yang-Baxterization of the Q -system. Thus one may say that the degree of the expansion expresses the degree of a Yang-Baxterization. Using $\{b_n^{(a)}(0)\}$, we can calculate the free energy (4.8) and the specific heat $C = -T \frac{\partial^2 f}{\partial T^2}$. We have plotted the high temperature expansion of the specific heat for $sl(3)$ in figure 4. For large T , this agrees with the result from another NLIE by Fujii and Klümper (Fig.4 in [18]). This indicates the validity of our new NLIE (4.7). We note that the high temperature expansion for the $SU(n)$ Heisenberg model is briefly reported in the letter [31] based on a completely different method.




Figure 4: Temperature dependence of the high temperature expansion of the specific heat of order 8 for $sl(3)$, $J = 1$, $\mu_1 = h$, $\mu_2 = 0$, $\mu_3 = -h$ case ($h = 0$: smooth line; $h = 2$: broken line; $h = 4$: dotted line.)

6 Concluding remarks

In this paper, we have derived a system of NLIE with a *finite* number of unknown functions, which describes thermodynamics of the $sl(r+1)$ Uimin-Sutherland model. This type of NLIE for $sl(r+1)$ of *arbitrary* rank r is derived for the first time. In particular for $r=1$, our new NLIE (4.7) reduces to Takahashi's NLIE [19] for the XXX spin chain. A relation between our new NLIE (4.6), (4.7) and the traditional TBA equations (3.11), (3.13), (3.16) which are also derived from the T -system is clarified. The high temperature expansion of the free energy is discussed, in which a solution of the T -system is given in terms of solutions of the Q -system. We expect that we can extend these results to other algebras by using the T -systems in [21, 32, 34, 35, 36, 37, 23].

In [38], fugacity expansion formulae of the free energy for the XXX spin chain up to an infinite order is derived from the string center equation, and Takahashi's NLIE [19] is rederived from this formulae. This work may be viewed in part as a kind of Yang-Baxterization of the results in [39] where power series formulae related to a formal completeness of the Bethe ansatz are derived from the string center equation. We can also generally recover [40] the results in [39] from the Q -system. Starting from our new NLIE (4.7), we may derive expansion formulae similar to the ones in [38] for $sl(r+1)$ of arbitrary rank r , from which an idea toward a Yang-Baxterization of the results in [40] may be given since our new NLIE is based on the T -system which is a Yang-Baxterization of the Q -system.

There are different types of NLIE with finite numbers of unknown functions for algebras of arbitrary rank in rather different contexts [41, 42]. Their origin are related to Destri and de Vega's NLIE [43]. To derive analogous NLIE from the QTM method and to study the relation to our new NLIE deserve investigation.

Acknowledgments

The author is financially supported by Inoue Foundation for Science.

References

- [1] M. Takahashi, Thermodynamics of One-Dimensional Solvable models, (Cambridge University Press, Cambridge, 1999).
- [2] M. Takahashi, *Prog. Theor. Phys.* **46** (1971) 401-415.

- [3] M. Gaudin, *Phys. Rev. Lett.* **26** (1971) 1301.
- [4] G. Jüttner, A. Klümper and J. Suzuki, *Nucl. Phys.* **B512** (1998) 581-600.
- [5] M. Suzuki, *Phys. Rev.* **B31** (1985) 2957-2965.
- [6] M. Suzuki and M. Inoue, *Prog. Theor. Phys.* **78** (1987) 787-799.
- [7] T. Koma, *Prog. Theor. Phys.* **78** (1987) 1213-1218.
- [8] J. Suzuki, Y. Akutsu and M. Wadati, *J. Phys. Soc. Jpn.* **59** (1990) 2667-2680.
- [9] A. Klümper, *Ann. Physik* **1** (1992) 540-553.
- [10] A. Kuniba, K. Sakai and J. Suzuki, *Nucl. Phys.* **B525** (1998) 597-626.
- [11] K. Sakai and Z. Tsuboi, *Int. J. Mod. Phys.* **A15** (2000) 2329-2346; math-ph/9912014.
- [12] Z. Tsuboi, *Int. J. Mod. Phys.* **A17** (2002) 2351-2368; cond-mat/0108358.
- [13] M. Takahashi, M. Shiroishi and A. Klümper, *J. Phys. A: Math. Gen.* **34** (2001) L187-L194.
- [14] A. Klümper, *Z. Phys.* **B91** (1993) 507-519.
- [15] G. Jüttner and A. Klümper, *Euro. Phys. Lett.* **37** (1997) 335-340.
- [16] G. Jüttner, A. Klümper and J. Suzuki, *Nucl. Phys.* **B522** (1998) 471-502.
- [17] J. Suzuki, *J. Phys. A: Math. Gen.* **32** (1999) 2341-2359.
- [18] A. Fujii and A. Klümper, *Nucl. Phys.* **B546** (1999) 751-764.
- [19] M. Takahashi, in *Physics and Combinatorics*, eds. A. N. Kirillov and N. Liskova, (2001) 299-304 (World Scientific, Singapore); cond-mat/0010486.
- [20] A. Klümper and P. Pearce, *Physica* **A183** (1992) 304-350.
- [21] A. Kuniba, T. Nakanishi and J. Suzuki, *Int. J. Mod. Phys.* **A9** (1994) 5215-5266.
- [22] Z. Tsuboi, *Phys. Lett.* **B544** (2002) 222-230; math-ph/0209024.

- [23] Z. Tsuboi, *J. Phys. A: Math. Gen.* **32** (1999) 7175-7206.
- [24] G.V. Uimin, *JETP Lett.* **12** (1970) 225.
- [25] B. Sutherland, *Phys. Rev.* **B12** (1975) 3795-3805.
- [26] J. H. H. Perk and C. L. Schultz, *Phys. Lett.* **84A** (1981) 407-410.
- [27] N. Yu. Reshetikhin and P. B. Wiegmann, *Phys. Lett.* **B189** (1987) 125-131.
- [28] A. N. Kirillov and N. Yu. Reshetikhin, *J. Sov. Math.* **52** (1990) 3156-3164.
- [29] V. V. Bazhanov and N. Reshetikhin, *J. Phys. A Math. Gen.* **23** (1990) 1477-1492.
- [30] M. Shiroishi and M. Takahashi, *Phys. Rev. Lett.* **89** (2002) 117201.
- [31] N. Fukushima and Y. Kuramoto, *J. Phys. Soc. Jpn.* **71** (2002) 1238-1241.
- [32] A. Kuniba and J. Suzuki, *J. Phys. A: Math. Gen.* **28** (1995) 711-722.
- [33] J. Suzuki, *J. Phys. A: Math. Gen.* **33** (2000) 3507-3521.
- [34] Z. Tsuboi, *J. Phys. A: Math. Gen.* **30** (1997) 7975-7991.
- [35] Z. Tsuboi, *Physica* **A252** (1998) 565-585.
- [36] Z. Tsuboi, *J. Phys. A: Math. Gen.* **31** (1998) 5485-5498.
- [37] Z. Tsuboi, *Physica* **A267** (1999) 173-208.
- [38] G. Kato and M. Wadati, *J. Math. Phys.* **43** (2002) 5060-5078.
- [39] A. Kuniba and T. Nakanishi, *Progr. in Math.* **191** (2000) 185-216.
- [40] A. Kuniba, T. Nakanishi and Z. Tsuboi, *Commun. Math. Phys.* **227** (2002) 155-190; math.QA/0105145.
- [41] P. Zinn-Justin, *J. Phys. A: Math. Gen.* **31** (1998) 6747-6770.
- [42] P. Dorey, C. Dunning and R. Tateo, *J. Phys. A: Math. Gen.* **33** (2000) 8427-8441.
- [43] C. Destri and H. J. de Vega, *Phys. Rev. Lett.* **69** (1992) 2313-2317.